

Detection of *Phytophthora* species in Plantations in S. Italy by a PCR-based Diagnostic



S.O. Cacciola (Dipartimento S.En.Fi.Mi.Zo., University of Palermo, viale delle Scienze 2, 90129 Palermo, Italy)

G. M. de San Lio, Dipartimento di Agrochimica e Agrobiologia, Università Mediterranea di Reggio Calabria

G.E. Agosteo (Dipartimento di Agrochimica e Agrobiologia), Mediterranean University of Reggio Calabria, piazza S. Francesco di Sales 4, 89061 Gallina(RC), Italy)

M. Funaro (A.R.S.S.A Regione Calabria, Ce.D.A. n.16, contrada Mirelli, Catanzaro Lido (CZ), Italy)

D. E. Li. Cooke, V. Young, J. M. Duncan, Scottish Crop Research Institute, Invergowrie, Dundee, DD2 5DA, U

Introduction

Up to 5 *Phytophthora* species have been recorded on strawberry in Europe. *P. fragariae* and *P. cactorum*, which cause red stele and crown rot respectively, are the commonest and most important species. Crown rot has been recorded in Italy but red stele has not.

Plantations around Lamezia Terme in S. Italy at the end of 2001 were affected by root rot, collapse and typical symptoms of crown rot

but several attempts to isolate *Phytophthora* from them failed. Samples of two main stocks of Camarosa, one from California and the other Belarus, were tested with a polymerase chain reaction diagnostic (PCRD) for *Phytophthora*. The basis of the test is explained in the poster and the results from S. Italy are given in the poster below



Strawberry red stele disease



Strawberry crown rot

Materials and Methods

Samples

The samples came from a single large plantation near Lamezia Terme. The area had been planted with Camarosa with nursery stock from California and Belarus. Patches of poor growth and general plant collapse were found throughout the field shortly after planting. Some affected plants had discoloured crowns typical of crown rot and most plants evidently had failed to make any growth after planting, suggesting that they were affected at time of planting. Samples of crowns and roots were inserted into apples to try and recover live *Phytophthora* cultures and isolations were attempted onto selective medium.

in the Internally Transcribed Spacer (ITS) regions of Ribosomal DNA (Bonants et al., 1995). The PCRD uses nested, or two-round, PCR in which DNA of any member of the Peronosporales present in the sample, e.g. a *Phytophthora*, *Pythium* or downy mildew, is amplified in the first round but DNA from other microorganism and plants is not amplified. Thereafter, in the second round PCR primers specific for particular *Phytophthora* species and internal to the amplicon from the first round can be used to test for particular species of interest, in this case *P. fragariae* and *P. cactorum*. Roots and crowns were frozen in liquid nitrogen and then ground very finely in a mortar and pestle before DNA extraction and purification.

PCR specificity and sensitivity


PCR testing

The polymerase chain reaction diagnostic (PCRD) is based on DNA sequence differences

Primers for nested PCR detection of *Phytophthora*

	Sequence	Name	Annealing temperature
1st round primers for all Peronosporales			
forward	5'GAGGGACTTTGGGGTAATCA 3'	DC6	Tm = 62oC
reverse	5' TCCTCCGCTATTGATATGC 3'	ITS4	
2nd round primers for specific for various <i>Phytophthora</i> species			
<i>P. fragariae</i>	forward 5'ACTAGTGGGGCCTGTCT 3'	DC1	Tm = 62oC
<i>P. fragariae/cambivora/cinnamomi</i> *	reverse 5'CGCCGACTGGCCACACAG 3'	DC5	

* Different forward primers are required to detect *P. cambivora* and *P. cinnamomi*



Pictures from Lamezia Terme

1. Plants being planted
2. Wilted plants
3. plants with rotted roots
4. Rotting in the crowns

Results

Apple baits and selective medium


No *Phytophthora* was recovered from the apple baits or the selective agar, perhaps because the material was very badly rotted by the time it was available for testing.

Nested PCR

Phytophthora was detected in several of the samples by PCR. Both *P. fragariae* and *P. cactorum*, were detected in samples from the stocks from Belarus, whereas only *P. cactorum* was detected in one sample from California. The same PCR techniques have been applied in several surveys in different countries. Typical results obtained on testing strawberry nursery stock with the PCRD are shown in the panel at left. The top panel shows the results after the first round of nested PCR using the Peronosporales primers DC6 and ITS4. There are many positives possibly because of *Pythium* and relatives being present in roots. The middle panel shows the second round of nested PCR using the DC1 and DC5 primers (see above), are specific for *P. fragariae* - four of the sample are positive. The bottom panel shows the same samples but this time, the primers specific for *P. cactorum* have been used (ADF1 and ADR in the panel above). There are six positives and one sample with an abnormal signal (highlighted with a red box), a very unusual occurrence. One sample (number 10) is positive for both pathogens.

Improved version of PCRD

The top panel In recent versions of the PCRD the DC6/ITS4 primers have been replaced with a set of primers that are specific for all *Phytophthora* species. These primers have increased sensitivity and eliminated many of the positives previously obtained in the first round of



Conclusions

P. cactorum was detected by the PCRD in strawberry plants showing poor growth and typical crown rot symptoms in the plantations at Lamezia Terme in Italy, although the pathogen was not isolated by either bait plants or selective media, mostly probably because of the poor condition of the plants at sampling. *P. cactorum* was detected in nursery plants from both California and Belarus; *P. fragariae* was also detected in plants from Belarus although there was no obvious symptoms of red core. These results and those of much

Acknowledgements

The authors acknowledge the joint assistance of the British Council and the Italian Academic Ministero dell'Istruzione dell'Università e della Ricerca which made this collaboration possible. We also wish to thank the Scottish Executive Environment and Rural Affairs Department, Horticultural Development Council and the EU Standards Testing and